

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

**EP 1 020 527 A1**

(12)

**EUROPEAN PATENT APPLICATION**

published in accordance with Art. 158(3) EPC

(43) Date of publication:

19.07.2000 Bulletin 2000/29

(51) Int. Cl.7: **C12N 15/55**

(21) Application number: 99933237.2

(86) International application number:

PCT/JP99/04167

(22) Date of filing: 03.08.1999

(87) International publication number:

WO 00/08176 (17.02.2000 Gazette 2000/07)

(84) Designated Contracting States:

AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU  
MC NL PT SE

(72) Inventors:

- HAMADA, Kazuyuki,  
Japan Tobacco Inc.  
Shizuoka 438-802 (JP)
- NAKAKIDO, Fumio,  
Japan Tobacco Inc.  
Shizuoka 438-0802 (JP)

(30) Priority: 04.08.1998 JP 22006098

(71) Applicant: Japan Tobacco Inc.

Tokyo 105-8422 (JP)

(74) Representative:

Reinhard - Skuhra - Weise & Partner  
Postfach 44 01 51  
80750 München (DE)

(54) **MUTATED BARNASE GENE AND PLANT TRANSFORMED BY THE SAME**

(57) A male sterile plant which is free from any undesirable character is constructed by expressing a barnase gene alone specifically in the anther. The barnase gene, which has been depressed by mutagenesis while sustaining its activity, is transferred into a plant and then expressed specifically in the anther to thereby efficiently give the aimed male sterile transformant.

EP 1 020 527 A1

## Description

[0001] The present application claims priority from Japanese Patent Application No.: Hei 10-220060, the disclosure of which is incorporated herein by reference.

## FIELD OF THE INVENTION

[0002] This invention relates to a mutant barnase gene which makes it possible to efficiently yield a male sterile transgenic plant when expressed in a specific site of a plant, in particular, anther-specifically. The present invention further relates to a recombinant vector capable of expressing the mutant barnase gene of the invention in host cells, a plant transformed by this vector and a method for constructing a transgenic plant.

## PRIOR ART

[0003] Barnase is an RNase originating in *Bacillus amyloliquefaciens* (S. Nishimura and M. Nomura, Biochem. Biophys. Acta 30, 430-431:1958; R.W. Hartley, J. Mol. Biol., 202, 913-915:1988). This enzyme has 110 amino acid residues and hydrolyzes RNA. When expressed in cells, this enzyme inhibits the functions of the cells as a result of its potent RNase activity and thus causes cell death in many cases. By using this characteristic, it is therefore expected that the function of the specific site can be selectively controlled by expressing the barnase gene in a specific site of a plant.

[0004] PCT International Publication WO89/10396 discloses a technique whereby a male sterile plant is obtained by constructing a male sterility gene by ligating the above-described barnase gene to the downstream of an anther tapetal cell-specific expression promoter and introducing the thus obtained gene into a plant. Male sterilization techniques are of great value in efficiently developing an F1 hybrid variety.

[0005] When the barnase gene is employed as a male sterility gene, however, it is frequently observed that resulting male sterile transgenic plants exhibit unfavorable characteristics. PCT International Publication WO96/26283 refers to this problem in rice. It is also reported that similar phenomena are observed not only in rice but in lettuce (Scientia Horticulturae 55, 125-139:1993; Arlette Reymaerts, Hilde Van de Wiele, Greta De Sutter, Jan Janssens: Engineered genes for fertility control and their application in hybrid seed production). According to this report, a plant with depressed activity was constructed by introducing a male sterility gene comprising a tobacco anther-specific promoter (TA29) and a barnase gene into lettuce.

[0006] Although reasons for these phenomena have not been accurately clarified so far, it is assumed that the so-called "position effect" of the site in which the gene is transferred may account for the mechanism (PCT International Publication WO96/26283). More specifically, a desired male sterile plant should be constructed if the male sterility gene is expressed exclusively in the target site (i.e., anthers). However, there is a possibility that the barnase might be expressed also in tissues other than anthers although in a very small dose under the action of expression regulators (for example, an endogenous enhancer) existing in the vicinity of the gene transfer site. In such a case, the unfavorable characteristics as described above can appear because barnase has a strong enzymatic activity.

[0007] To overcome this problem, the method disclosed in PCT International Publication WO96/26283 exploits the character of cauliflower mosaic virus 35S promoter (hereinafter referred to as CaMV35S promoter) of being expressed potently in tissues other than anthers. Namely, barstar, i.e., an inhibitory protein against barnase is employed therein. The barstar gene ligated to a CaMV35S promoter is transferred into a plant simultaneously with a barnase gene and then the barstar gene is constitutively expressed in tissues other than the anthers, thereby eliminating the effects of the barnase in tissues other than the anthers. However, it is necessary in this method to transfer not only a barnase gene but also a barstar gene, and hence, the problem is that the application of this technique to the breeding of F1 varieties of rice or corn seeds may give rise to a gene silencing. "Gene silencing" is a phenomenon wherein the expression of a gene is inhibited when plural copies of the foreign gene are introduced. It is known that this problem frequently occurs when the expression of a foreign gene is effected by a 35S promoter, though the detailed mechanism thereof has not been clarified yet (R.B. Flavell, Proc. Natl. Acad. Sci. USA 91, 3490-3496:1994; J. Finnegan, Bio. Technology 12, 883-888:1994; M.A. Matzke and A.J.M. Matzke, Plant Physiol., 107, 679-685:1995). Since, in rice and corn, a barstar gene is used as a "fertility-restoring gene" in the pollen parent (father) so as to allow the pollen in the F1 generation to restore the fertility (C. Mariani, et al., Nature 357, 384-387:1992), if an MS plant (mother) is constructed by the method with the use of a barstar gene as reported in WO96/26283, the F1 plant has plural copies of the barstar gene. In such a case, there is a possibility that the expression of the gene may be inhibited due to the gene silencing.

## SUMMARY OF THE INVENTION

[0008] The present invention provides a method for constructing a male sterile plant by using a barnase gene with-

out resort to a barstar gene.

[0009] The present invention further provides a mutant barnase gene to be used in the above method and a process for producing the same.

## 5 DETAILED DESCRIPTION OF THE INVENTION

[0010] In the present invention, the DNA sequence of barnase gene (R.W. Hartley, J. Mol. Biol. 202, 913-915:1988) is mutated at least in part and then the thus obtained mutant barnase gene is anther-specifically expressed in a plant so as to make the plant substantially male sterile without any substantially disadvantageous effect on the tissues other than anthers.

[0011] The mutation can be performed by a known method such as site-specific mutagenesis, deletion of a fragment by using restriction enzymes or the low fidelity PCR method. Among all, it is preferred to use the low fidelity PCR method. This method is described in detail in D. Leung, E. Chen and D. Goedda, Technique 1, 11-15:1989; Y.Z. Xiaoping and R.H. Ebright, Nucleic Acid Res. 19, 6052:1991; G.C. Rice et al., Proc. Natl. Acad. Sci. USA 89, 5467-5471: 15 1992 and the contents of these documents are incorporated herein by reference. By using this technique, PCR is carried out under such conditions as to induce some errors during the amplification reaction. Thus, random mutations can be introduced into the target DNA fragment.

[0012] Primers to be used in the low fidelity PCR method in the present invention are selected as in the conventional PCR method. It is preferred that each of these primers has a similar number of nucleotides as in the conventional PCR method.

[0013] The present inventors performed PCR by using a DNA containing the sequence represented by SEQ ID NO:1 as a template with the use of a combination of primer 1 (5'-CGTTCGGCTC GATGGTACCG GTTATCAACA CGTTTGA-3'; SEQ ID NO: 6) and primer 2 (5'-CCTCTAGATT ATCTGATTTT TGTAAGGTC TGATAATG-3'; SEQ ID NO:7) under such conditions as to induce errors. Thus, a mutant barnase gene having the DNA sequence represented by SEQ ID NO:3 was isolated. The sequence represented by SEQ ID NO:1 employed herein is the sequence of the coding region of the barnase gene contained in the known plasmid pVE108 (WO92/ 13956). This sequence corresponds to the wild type barnase gene from which the unnecessary moiety corresponding to the secretion signal in the N-terminal side has been deleted.

[0014] It is also possible to acquire different mutant barnase genes by a similar method.

[0015] The PCR amplification product is cloned in a host (for example, *Escherichia coli*) and clones containing a mutant barnase gene is isolated. In this cloning, clones having a mutant barnase gene may be screened by assaying the RNase activity expressed by the gene. However, it is advantageous to screen the clone by the method consisting of the following two steps by taking advantage of the fact that the RNase activity of barnase affects the growth of *E. coli*.

[0016] In the first step, a plasmid is prepared by using the mutant barnase gene obtained above and then *E. coli* is transformed thereby. The growth of the *E. coli* transformant thus obtained will be inhibited by the activity of the mutant barnase. Based on this fact, a colony growing slowly (i.e., a small colony), compared with *E. coli* having a control vector free from any barnase gene, is selected. It is expected that the thus selected *E. coli* strain will contain the mutant barnase gene. To confirm that the inhibition of the growth of *E. coli* is due to the expression of the mutant barnase gene, the second step is then carried out.

[0017] In the second step, a barstar gene is employed. As described above, barstar is an inhibitory protein against barnase. In *E. coli* which expresses barstar, the enzymatic activity of barnase is inhibited and thus the degradation of mRNA, which otherwise takes place in the presence of barnase, can be inhibited. Therefore, *E. coli* which expresses a barstar gene is transformed with a barnase gene, the growth will not be inhibited. Accordingly, it is expected that the thus obtained *E. coli* transformant will produce little difference in growth rate from the transformant constructed by the control plasmid which is free from any barnase gene and, therefore, these transformants will produce little difference in colony size too. Based on this principle, a plasmid is prepared from the *E. coli* strain selected in the first step. The plasmid is then used to transform another *E. coli* strain, in which the barstar gene is constitutively expressed. Thus, mutant barnase gene-containing colonies having almost the same size as the one of the *E. coli* transformed by the control plasmid are selected. The *E. coli* with the constitutive expression of the barstar gene to be used herein can be prepared by, for example, the method as will be described in Example 1 hereinafter.

[0018] The DNA sequence of the thus obtained mutant barnase gene may be analyzed by a conventional method, if necessary, to thereby examine the mutation in detail.

[0019] A preferred example of the mutant barnase gene of the present invention is one which has the DNA sequence represented by SEQ ID NO:3. Compared with the nucleotide sequence of SEQ ID NO:1 coding for the wild type activity, the nucleotide sequence encoding this gene has an insertion of "T" at the 15-position from A in the initiation codon ATG and a deletion of "A" at the 333-position. According to the normal translation manner, translation will be initiated from ATG at the 1-position of this DNA sequence represented by SEQ ID NO:3, and terminated at the ninth codon which has been converted to a termination codon as a result of the insertion of the "T" at the 15-position. It is

unlikely that the thus formed oligopeptide consisting of 8 amino acids has the barnase activity. Further, there is no other ATG or GTG codon from which the translation can be initiated in the correct frame in the vicinity thereof. However, the fact that the growth rate of *E. coli* was restored by the barstar protein strongly suggests that the translation of the barnase protein from the gene of SEQ ID NO:3 proceeded in the correct frame. This is seemingly due to a phenomenon called "frame shift re-coding" whereby the ribosome controls the translation while automatically shifting the reading frame during the course of translation into the protein. This phenomenon, which has been reported in several genes of viruses, *E. coli* and animals, generally depends exclusively on an individual nucleotide sequence and neither any special protein nor translation device is required. The frame shifting efficiency during the translation varies from gene to gene, i.e., ranging from several % to 50%. In the case of the mutant barnase gene, it is considered that the efficiency of the shifting of the reading frame during the translation is not high and thus the translation product in the correct frame is formed in a smaller amount, thereby lowering the activity of the gene.

[0020] Moreover, there is a high possibility that other mutant barnase genes of which the translation efficiency is lowered due to the shifting of the reading frame, as in the case of the DNA sequence represented by SEQ ID NO:3, or some other reasons can be obtained by mutating the barnase gene by, for example, the low fidelity PCR method. Furthermore, there is a high possibility that a gene having a DNA sequence derived from the DNA sequence represented by SEQ ID NO:3 by substitution, deletion, insertion or addition of one or more nucleotides will be a mutant barnase gene which provides a lowered translation efficiency due to the shifting of the reading frame as a result of the insertion of T at the 15-position from A in the initiation codon ATG and/or the deletion of A at the 333-position, as in the DNA sequence represented by SEQ ID NO:3. It is considered that each of these genes, likewise the gene of SEQ ID NO:3, is capable of making a plant substantially male sterile when it is anther-specifically expressed in the plant. Accordingly, these genes are also included in the scope of the present invention similar to the gene of SEQ ID NO:3.

[0021] The plant which is to be male sterilized by the mutant barnase gene is not particularly limited, so long as the gene can be transferred into the plant to provide a male sterilized plant. Examples of the plants include rice, corn, tobacco, lettuce and Brassica. Among all, rice and corn are preferred.

[0022] In order to male sterilize a plant, the mutant barnase gene is expressed specifically in the anthers of the plant to inhibit the anther functions by the RNase activity of barnase. Such an anther-specific expression can be achieved by using the method described in WO92/13957. In brief, the mutant barnase gene is ligated to the downstream of an anther-specific promoter and then incorporated into plant cells with the use of an expression vector.

[0023] The incorporation may be performed by using the Agrobacterium method, the electroporation method, the particle gun method, etc.

[0024] From the transformed plant cells, a complete plant can be formed by the regeneration from a callus of the transformed plant cells in accordance with, for example, the method reported in Y. Hiei et al., Plant J. 6, 271-282:1994.

[0025] The male sterility of the thus constructed plant can be confined by the inability of the plant to fertilize unless it is pollinated from another fertilizable plant.

#### Example 1: Preparation of mutant barnase gene

##### Low fidelity PCR

[0026] Primer 1 (CGTTCGGCTC GATGGTACCG GTTATCAACA CGTTTGA: SEQ ID NO: 6) and Primer 2 (CCTCTAGATT ATCTGATTTT TGTAAGGTC TGATAATG: SEQ ID NO:7) were synthesized by a DNA synthesizer (manufactured by Applied Biosystems) in accordance with the method described in S.L. Beaucage et al., Tetrahedron Lett., 22, 1859-1862:1982. Known plasmid pVE108 (WO92/13956) was used as a template and PCR was carried out with the use of the combination of Primers 1 and 2. SEQ ID NO:1 shows the sequence of the coding region of the barnase gene which is contained in pVE108. The reaction was performed for 50 cycles with each cycle consisting of 1 minute at 94°C, 1 minute at 57°C and 1 minute at 72°C in 10 mM Tris-HCl (pH 9.5), 50 mM KCl, 2 mM MgCl<sub>2</sub>, and 1 mM each of dNTP, 10 ng of the template DNA and 0.5 U of Taq DNA polymerase.

[0027] After the completion of the reaction, the amplification product was separated by electrophoreses on agarose gel (2% SeaKem GTG agarose, 1xTAE) and purified by the DEAE-cellulose method (M. Muramatsu, Labo-manual Idenshikogaku (Gene Engineering Labo-Manual), Maruzen, pp.111:1988). By using the purified product thus obtained as a template, PCR was performed again under the conditions as defined above.

##### Ligation of mutated barnase gene fragment into plasmid vector

[0028] The reaction product from the above step was digested with SacI and XbaI in a conventional manner and then purified by agarose gel electrophoresis to give an "insertion fragment". A plasmid to be used for transferring this insertion fragment into *E. coli* can be appropriately selected. For example, use may be made of plasmid pHM1 therefor. The plasmid pHM1 was constructed as follows. Plasmid pBR322 was cleaved at the EcoRI site and blunt-ended by

using T4 DNA polymerase (manufactured by Takara Shuzo). Separately, lacZ expression cassette (322 bp) was excised from pUC18 with PvuII and the ends thereof were blunted. The cassette was integrated into the restriction site of the blunt-ended plasmid pBR 322 to give plasmid pHM1.

5 [0029] Plasmid pHM1 was digested with SacI and XbaI followed by dephosphorylation with calf intestine alkaline phosphatase (manufactured by Takara Shuzo) to give a restriction enzyme-treated plasmid fragment. Into this restriction enzyme-treated plasmid pHM1, the above "insertion fragment" containing a barnase gene was ligated by using Takara Ligation Kit ver.1 (manufactured by Takara Shuzo).

#### Introduction into *E. coli* and selection of barnase-active clone

10 [0030] The *E. coli* having the barnase gene introduced thereto may be selected by, for example, the following method. Since mRNA is degraded in cells by the barnase activity, the synthesis of protein is suppressed and, in its turn, the growth of *E. coli* is inhibited. Thus, the *E. coli* transformed by the barnase gene can be selected by taking advantage of the fact that the desired transformant colony has a smaller size than the *E. coli* colony transformed by the control plasmid free from the mutant barnase gene. When the wild type barnase or a mutant barnase still maintaining the activity comparable thereto is integrated into pHM1, *E. coli* cannot form any colony. Thus, clones having a sufficiently weakened barnase activity can be exclusively selected.

15 [0031] Thus, the plasmid having the mutant barnase gene ligated thereto was precipitated from ethanol and then introduced into *E. coli* LE392 strain by the electroporation method with the use of GenePulser (BioRad) in accordance with the manufacture's instruction. Also, the control plasmid free from mutant barnase was introduced into the *E. coli* LE392 strain in the same manner.

20 [0032] Next, these *E. coli* transformants were plated onto an LB agar medium containing tetracycline (25 µg/ml) and incubated at room temperature (25°C) for 72 hours. Then colonies of a smaller size than the colony of *E. coli* having the control plasmid pHM1 introduced thereto were screened as colonies of the *E. coli* transformants having the mutant barnase transferred thereto (Table 1, A).

#### Selection of mutant barnase clone by barstar gene

30 [0033] As described above, barstar is an enzyme acting as an antagonist to barnase. In *E. coli* wherein barstar is expressed, the enzymatic activity of barnase is inhibited and thus the degradation of mRNA, which otherwise takes place in the presence of barnase, can be inhibited. Therefore, if *E. coli* expressing barstar is transformed by a plasmid containing a barnase gene, the growth thereof will not be inhibited. Accordingly, it is expected that the thus obtained *E. coli* transformant will show little difference in growth rate from the transformant constructed by using the control plasmid free from a barnase gene and, therefore, these transformants will show little difference in colony size too.

35 [0034] An *E. coli* transformant which expressed barstar was constructed in the following manner. A barstar gene described in R.W. Hartley, J. Mol. Biol. 202, 913-915:1998 was excised with HindIII and XbaI and ligated in frame to a tac promoter (de Boer et al., Proc. Natl. Acad. Sci. USA 80, 21-25:1983). The gene, together with a chloramphenicol tolerance gene (N.K. Alton and D. Vapnek, Nature 282, 864-869:1979), was ligated into the defective transposon which lacked the transferase in a vector described in Herrero et al., J. Bacteriol. 172, 6557-6567:1990. Subsequently, the thus obtained plasmid was introduced into *E. coli* MC1061 strain thereby to transfer the transposon, which contained the barstar gene cassette, into the chromosome of the *E. coli*. It was confirmed based on the chloramphenicol tolerance that this *E. coli* was able to maintain the barstar gene cassette in a stable state. Since the *E. coli* MC1061 strain lacked the lacI gene, the tac promoter was continuously active and thus the barstar gene was constitutively expressed.

40 [0035] The plasmid containing the mutant barnase gene as the insertion fragment and the control plasmid free from the same were each introduced into the above *E. coli* transformant by the electroporation method. Then the *E. coli* transformants were plated onto an LB agar medium containing IPTG (1mM) and tetracycline (25 µg/ml) and incubated at 25°C for 72 hours. Among the colonies of the *E. coli* transformed by the mutant barnase gene, a colony being comparable in size to the colony of the *E. coli* having the control plasmid pHM1, which had been simultaneously plate-cultured, was screened and named "#4-31" (Table 1, B).

Table 1

Screened clone and colony size (mm)		
	Screened clone	pHM1 (control)
A: <i>E. coli</i> LE392* <sup>1</sup>	1.77 ± 0.33	2.65 ± 0.47
B: <i>E. coli</i> with barstar gene expression* <sup>2</sup>	1.39 ± 0.10	1.55 ± 0.08

\*1: Colony size after incubation on the LB agar plate containing 25 µg/mL of tetracycline at 28°C for 48 hours.

\*2: Colony size after incubation on the LB agar plate containing 25 µg/mL of tetracycline at 28°C for 24 hours.

#### Example 2: Determination of nucleotide sequence of mutant barnase gene

[0036] From the *E. coli* screened in Example 1, a cloned mutant barnase gene appropriate for the object of the present invention was prepared. First, the clone #4-31 screened in Example 1 was propagated. Next, the mutant barnase gene fragment in the clone #4-31 was excised with KpnI and XbaI and ligated into the KpnI, XbaI-site of pUC119 having been cleaved similarly with KpnI and XbaI. After amplifying the obtained plasmid by using *E. coli*, a reaction was performed by the cycle sequence method with the use of Taq polymerase (Taq Dye Terminator Cycle Sequencing Kit, manufactured by Applied Biosystems Inc.) in accordance with the manufacturer's protocol. Subsequently, the sequence was analyzed with a DNA Sequencer (Model 373A, manufactured by Applied Biosystems Inc.), thereby yielding the sequence represented by SEQ ID NO:3. Compared with the DNA sequence of the wild type barnase gene, this sequence had an insertion of T at the 15-position from A of the initiation codon ATG and a deletion of A at the 333-position.

#### Example 3: Construction of male sterile rice by using attenuated mutant barnase gene

[0037] The attenuated mutant barnase gene integrated into pUC119 as described above was cleaved with restriction enzymes XbaI and KpnI and thus the plasmid vector pTS431 represented by SEQ ID NO:5 was constructed. Although this plasmid pTS431 differs from the known plasmid pVE108 (PCT International Publication WO92/13956) in the following points, it is substantially equivalent thereto except for the moieties of the anther-specific promoter and the mutant barnase.

(1) In the plasmid vector pTS431 according to the present invention, the barnase gene (SEQ ID NO:1) in the known plasmid pVE108 has been converted into the mutant barnase gene (SEQ ID NO:3).

(2) A tobacco-origin anther specific promoter is used in the known plasmid pVE108, while the anther specific promoter of rice E1 gene (PCT International Publication WO92/13956) is used in the plasmid vector pTS431 according to the present invention.

(3) In the present invention, a 35S3 promoter (EP 0344029) of 1376 bp, which is not used in the plasmid pVE108, is employed upstream of the barnase gene.

(4) In the present invention, a sequence originating in the downstream of Agrobacterium T-DNA gene 7 excised from pJD884 (PCT International Publication WO93/09218) is employed whereas such a sequence is not used in the plasmid pVE108.

(5) In the present invention, the region corresponding to lacZ has been deleted from the pUC19-origin moiety. In addition a plasmid (pTS172) having the wild type barnase gene integrated therein is represented by SEQ ID NO:4.

[0038] The fragment of about 4.5 kbp was excised with a restriction enzyme EcoRI respectively from pTS431 (a plasmid having the mutant barnase gene transferred therein; SEQ ID NO:5) and pTS172 (a plasmid having the barnase gene transferred therein; SEQ ID NO:4) and inserted into the EcoRI-site of the intermediate vector pSB11 (T. Komari et al., Plant J. 10(1), 165-174:1996), and further, the T-DNA region thereof was integrated into the acceptor vector pSB1 (T. Komari et al., Plant J. 10(1), 165-174:1996) by way of homologous recombination. By using *Agrobacterium tumefaciens* LBA4404 having the obtained recombinant plasmids (pSB1431 and pSB1172 respectively), rice (variety: Asanohikari) was transformed. While the transformation was performed basically in accordance with the method of Hiei et al. (Plant J. 6(2), 271-282:1994), phosphinothricine (concentration: 10 mg/L) was employed for screening transformants since the thus constructed male sterility genes contained bar gene encoding phosphinothricine acetyl transferase as a selective marker. Phosphinothricine facilitates selection of calluses having the gene transferred therein.

[0039] A comparison of the rice transformant obtained with the use of the wild type barnase gene with the one obtained with the use of the mutant barnase gene indicates that the transfer of the mutant barnase gene gave rise to a remarkable improvement in the transformation efficiency and an increase in the rate of morphologically normal male sterility transformants, as shown in Table 2.

Table 2

Transformation efficiency				
	No. of infected calluses	No. of regenerated calluses	No. of PCR-positive lines* <sup>1</sup>	No. of morphologically normal male sterile lines
pSB1172 (control)	2838	83	52/83	9/52(17.3%)
pSB1431 (present invention)	787	69	43/45* <sup>2</sup>	27/28* <sup>3</sup> (96.4%)

\*1: detection of barnase gene fraction by PCR.

\*2: examined only 45 lines among 69.

\*3: examined only 28 lines among 43.

#### Effects of the Invention

[0040] In the present invention, a barnase gene having a weakened effect is constructed via mutation. A male sterility gene comprising the mutant barnase gene can be introduced into a plant to successfully produce a male sterile plant free from any unfavorable characteristic at a high efficiency with the use of a single gene without resort to the use of a barstar gene.

## SEQUENCE LISTING

5 <110> Japan Tobacco Inc.

<120> MUTANT BARNASE GENE AND TRANSGENIC PLANT TRANSFORMED BY SAID GENE

10 <130> 980687

15 <160> 7

<210> 1

20 <211> 343

<212> DNA

<213> Bacillus amyloliquefaciens

25 <220>

<221> NAME/KEY: CDS

30 <222> 1..336

<400> 1

35 atg gta ccg gtt atc aac acg ttt gac ggg gtt gcg gat tat ctt cag 48  
Met Val Pro Val Ile Asn Thr Phe Asp Gly Val Ala Asp Tyr Leu Gln  
1 5 10 15

40 aca tat cat aag cta cct gat aat tac att aca aaa tca gaa gca caa 96  
Thr Tyr His Lys Leu Pro Asp Asn Tyr Ile Thr Lys Ser Glu Ala Gln  
20 25 30

45 gcc ctc ggc tgg gtg gca tca aaa ggg aac ctt gca gac gtc gct ccg 144  
Ala Leu Gly Trp Val Ala Ser Lys Gly Asn Leu Ala Asp Val Ala Pro  
35 40 45

50 ggg aaa agc atc ggc gga gac atc ttc tca aac agg gaa ggc aaa ctc 192  
Gly Lys Ser Ile Gly Gly Asp Ile Phe Ser Asn Arg Glu Gly Lys Leu

55



EP 1 020 527 A1

50 55 60

5 cgc ggc aaa agc gga cga aca tgg cgt gaa gcg gat att aac tat aca 240  
Pro Gly Lys Ser Gly Arg Thr Trp Arg Glu Ala Asp Ile Asn Tyr Thr

65 70 75 80

10 tca ggc ttc aga aat tca gac cgg att ctt tac tca agc gac tgg ctg 288  
Ser Gly Phe Arg Asn Ser Asp Arg Ile Leu Tyr Ser Ser Asp Trp Leu

85 90 95

15 att tac aaa aca acg gac cat tat cag acc ttt aca aaa atc aga taa 336  
Ile Tyr Lys Thr Thr Asp His Tyr Gln Thr Phe Thr Lys Ile Arg

100 105 110

20 ggtaacc 343

<210> 2

25 <211> 112

<212> PRT

30 <213> Bacillus amyloliquefaciens

<400> 2

35 Met Val Pro Val Ile Asn Thr Phe Asp Gly Val Ala Asp Tyr Leu Gln

1 5 10 15

Thr Tyr His Lys Leu Pro Asp Asn Tyr Ile Thr Lys Ser Glu Ala Gln

40 20 25 30

Ala Leu Gly Trp Val Ala Ser Lys Gly Asn Leu Ala Asp Val Ala Pro

35 40 45

45 Gly Lys Ser Ile Gly Gly Asp Ile Phe Ser Asn Arg Glu Gly Lys Leu

50 55 60

Pro Gly Lys Ser Gly Arg Thr Trp Arg Glu Ala Asp Ile Asn Tyr Thr

50 65 70 75 80

Ser Gly Phe Arg Asn Ser Asp Arg Ile Leu Tyr Ser Ser Asp Trp Leu

55

EP 1 020 527 A1

		85		90		95	
5	Ile	Tyr	Lys	Thr	Thr	Asp	His
				Tyr	Gln	Thr	Phe
				Thr	Lys	Ile	Arg
		100		105		110	

10 <210> 3

<211> 342

<212> DNA

15 <213>

<400> 3

20 atggtaccgg ttattcaaca cgtttgacgg ggttgcggat tatcttcaga catatcataa 60

gctacctgat aattacatta caaaatcaga agcacaagcc ctccgctggg tggcatcaaa 120

agggaacctt gcagacgtcg ctccggggaa aagcatcggc ggagacatct tctcaaacag 180

25 ggaaggcaaa ctcccgggca aaagcggacg aacatggcgt gaagcggata ttaactatac 240

atcaggcttc agaaattcag accggattct ttaactaagc gactggctga ttacaaaac 300

30 aacggaccat tatcagacct ttacaaaat cagtaactta ga 342

<210> 4

35 <211> 6548

<212> DNA

<213> Escherichia coli LE392

40 <220>

<223> Clone: pTS172

<400> 4

45 aattcaagct tgacgtcagg tggcactttt cggggaaatg tgcgcggaac ccctatttgt 60

ttatitttct aaatacattc aaatatgtat ccgctcatga gacaataacc ctgataaaag 120

50 cttcaataat attgaaaaag gaagagtatg agtattcaac atttccgtgt cgccttatt 180

cccttttttg cggcattttg ccttctctgtt ttgtctcacc cagaaacgt ggigaaagta 240

55

aaagatgctg aagatcagtt ggggtcacga gtgggttaca tcgaactgga tctcaacagc 300  
 5 ggtaagatcc ttgagagttt tcgccccgaa gaacgttttc caatgatgag cactttttaa 360  
 gtictgctat gtggcgcggt attatcccgt attgacgccg ggcaagagca actcggctgc 420  
 cgcatatact attctcagaa tgacttggtt gagtactcac cagtcacaga aaagcatctt 480  
 10 acggatggca tgacagtaag agaattatgc agtgcgtcca taacctgag tgataacact 540  
 gcggccaact tacttctgac aacgatcgga ggaccgaagg agctaaccgc tttttgcac 600  
 aacatggggg atcatgtaac tcgccttgat cgttgggaac cggagctgaa tgaagccata 660  
 15 ccaaacgacg agcgtgacac cacgatgcct gtagcaatgg caacaacgtt gcgcaaaacta 720  
 ttaactggcg aactacttac tctagcttcc cggcaacaat taatagactg gatggaggcg 780  
 gataaagtgt caggaccact tctgcgtcgc gcccttcggg ctggctggtt tattgctgat 840  
 20 aaatctggag ccggtgagcg tgggtctcgc ggtatcattg cagcactggg gccagatggt 900  
 aagccctccc gtatcgtagt tatctacacg acggggagtc aggcaactat ggatgaacga 960  
 aatagacaga tcgctgagat aggtgccica ctgattaagc attggttaact gtcagaccaa 1020  
 25 gtttactcat atatacttta gattgattta aaacttcatt ttttaattta aaggatctag 1080  
 gtgaagatcc tttttggctc gagtctcatg accaaaatcc cttaacgtga gtttctgtc 1140  
 cactgagcgt cagaccccggt agaaaagatc aaaggatctt cttgagatcc ttttttctg 1200  
 30 cgcgtaatct gctgcttgca aacaaaaaaa ccaccgtac cagcgggtgt ttttttgcg 1260  
 gatcaagagc taccaactct ttttcgaag gtaactggct tcagcagagc gcagatacca 1320  
 35 aatactgtcc ttctagtgtg gccgtagtta ggccaccact tcaagaactc tgtagcaccg 1380  
 cctacatacc tcgctctgct aatcctgtta ccagtggtg ctgccagtgg cgataagtgc 1440  
 tctcttaccg ggttgactc aagacgatag ttaccggata aggcgcagcg gtcgggctga 1500  
 40 acgggggggt cgtgcacaca gccagcttg gagcgaacga cctacaccga actgagatac 1560  
 ctacagcgtg agcattgaga aagcgccacg cttcccgaag ggagaaaggc ggacaggtat 1620  
 ccggtaaagc gcagggtcgg aacaggagag cgcacgaggg agcttcagg gggaaacgcc 1680  
 45 tggatatctt atagtcctgt cgggtttcgc cactctgac ttgagcgtc atttttgtga 1740  
 tgctcgtcag gggggcggag cctatggaaa aacgccagca acgcggcctt ttacggttc 1800  
 ctggcctttt gctggccttt tgctcacatg ttctttcctg cgttatcccc tgattctgtg 1860  
 50 gataaccgta ttaccgcctt tgagttagct gataccgctc gccgcagccg aacgaccgag 1920  
 cgcagcgagt cagttagcga ggaagcggaa gagcgcccaa tacgcaaacc gcctctcccc 1980

EP 1 020 527 A1

gcgcggttggc ctgatcagaa ttcataatgca cgtgttcccc atctagtaac atagatgaca 2040  
5 ccgcgcgcgga taatttatcc tagtttgcgc gctatatattt gttttctatc gcgtattaaa 2100  
tgtataattg cgggactcta atcataaaaa cccatctcat aaataacgic atgcattaca 2160  
tgttaattat tacatgcita acgtaattca acagaaatta tatgataatc atcgcaagac 2220  
10 cggaacagg attcaatctt aagaaacttt attgccaaat gtttgaacga tctgcttcgg 2280  
aggttacctt atctgatttt tgtaaaggic tgataatggt ccgttgtttt gtaaatcagc 2340  
cagtcgcttg agtaaagaat ccggtctgaa tttctgaagc ctgatgtata gttaatatcc 2400  
15 gcttcacgcc atgttcgtcc gcttttggcc gggagtittgc cttccctgtt tgagaagatg 2460  
tctccgccga tgcttttccc cggagcgcgc tctgcaaggt tcccttttga tgccaccag 2520  
ccgagggctt gtgcttctga ttttgtaatg taattatcag gtagcttatg atatgtctga 2580  
20 agataatccg caaccccgic aaacgtgttg ataaccgga ccatcgcgac ggcttgatgg 2640  
atctcttgct ggacaccggg atgctaggat gggttatcgt ggccggcgic cgtgtgtgic 2700  
25 ttttgtaggc gccggcgagc gcgggggcaa tgtggcaggt ggtcacggt gcaagcgtgc 2760  
gcaagtgaat gcaacaacca aggacggtca tggcgaaagc acctcacgcg tccaccgtct 2820  
acaggatgta gcagtagcac ggtgaagaa gtgttgccc gtccattagg tgcattctca 2880  
30 ccgttggcca gaacaggacc gticaacagt taggttgagt gtaggacttt tacgttggtta 2940  
atgtatggca aatagtagta aattttggcc ccattggtct ggctgagata gaacatattc 3000  
tggaagcct ctagcatatc tttttgaca gctaaacttt gcttcttgcc ttcttggtct 3060  
35 agcaatgacg ttgccatgt cgtggcaaac atctggttaag gtaactgtat tcgtttgttc 3120  
ccttcaacgg ctcaatcccc acaggccaag ctatcctttc cttggcagta taggctcctt 3180  
gagagattat actaccattt ttaagtgcit ataaagacga tgctctctaa ccagatcgat 3240  
40 cagaaacaca aagttttagc agcgtaatat cccacacaca tacacacacg aagctatgcc 3300  
tcctcatttt ccgagagatt ctgacagtga ccagaatgic agaatgccat ttcattggga 3360  
caagtcgac cacaagcttc ttggtggagg tcaagggtgt ctattattat tcgctttcta 3420  
45 ggaaattatt cagaattagt gccttttctc ataatcttc tctgagccga tgtggttttg 3480  
gatttcattg ttgggagcta tgcagttgcg gatattctgc tgtggaagaa caggaaacta 3540  
tctcgggggg tccttgctgg ggcaacattg atatggttcc tgttcgatgt agtagaatac 3600  
50 aatataattc cgctcctttg ccagattgcc attcttgcca tgcttgatgat cttcatttgg 3660  
tcaaatgccg caccactctt ggacaggat tagctttatt tcctgtggag atggtagaaa 3720

55

actcagctta cagaaatggc atttcacgta gtataacgca agacattagg tactaaaact 3780  
 5 caactaactg ttccgaatt tcagggcccc tccaaggatc ccagaaatca tcattctctga 3840  
 acatgccttc agagaaatgg cattgaccgt ccattacaaa ctaacgtaca ctgtatctgt 3900  
 tctttacgac attgcatgtg gaaaggatct gaagagattt ctcttggtac ataataatct 3960  
 10 actcctttgc tacgttaata agagatgtaa aaacatgcaa cagttccagt gccaacattg 4020  
 tccaaggatt gtgcaattct ttctggagcg ctaaaatiga ccagattaga cgcatcagaa 4080  
 tattgaattg cagagttagc caataatcct cataatgta atgtgctatt gttgttcaact 4140  
 15 actcaatata gttctggact aacaatcaga ttgtttatga tattaagggtg gttggatctc 4200  
 tattggtatt gtcggcgatt ggaagttctt gcagcttgac aagtctacta tatattggta 4260  
 ggtattccag ataaatatta aattttaata aaacaatcac acagaaggat ctgcggccgc 4320  
 20 tagcctaggc ccgggccac aaaaatctga gcttaacagc acagttgctc ctctcagagc 4380  
 agaatcgggt attcaacacc ctcatatcaa ctactacgtt gtgtataacg gtccacatgc 4440  
 cggtatatac gatgactggg gttgtacaaa ggcggaaca aacggcgctc ccggagttgc 4500  
 25 acacaagaaa ttggcacta ttacagaggc aagagcagca gctgacgcgt acacaacaag 4560  
 tcagcaaca gacaggttga acttcatccc caaaggagaa gctcaactca agcccaagag 4620  
 ctttgctaag gcctaacaa gccaccacaa gcaaaaagcc cactggctca cgctaggaac 4680  
 30 caaaggccc agcagtgatc cagcccaaaa agagatctcc ttgccccgg agattacaat 4740  
 ggacgatttc ctctatcttt acgatctagg aaggaagttc gaaggtaag gtgacgacac 4800  
 35 tatgttcacc actgataatg agaaggtag cctcttcaat ttcagaaaga atgctgaccc 4860  
 acagatggtt agagaggcct acgcagcagg tctcatcaag acgatctacc cgagtaacaa 4920  
 tctccaggag atcaaatacc ttccaagaa ggttaagat gcagtaaaa gattcaggac 4980  
 40 taattgcatc aagaacacag agaagacat atttctcaag atcagaagta ctattccagt 5040  
 atggacgatt caaggcttgc ttcataaacc aaggcaagta atagagattg gactctctaa 5100  
 aaaggtagtt cctactgaat ctaaggccat gcatggagtc taagattcaa atcgaggatc 5160  
 45 taacagaact cgccgtgaag actggcgaac agttcataca gactctttta cgactcaatg 5220  
 acaagaagaa aatcttcgtc aacatggtgg agcacgacac tctggtctac tcaaaaaatg 5280  
 tcaagatac agtctcagaa gaccaaaggc ctattgagac ttttcaacaa aggataattt 5340  
 50 cgggaaacct cctcggttc cattgccag ctatctgtca cttcatcgaa aggacagtag 5400  
 aaaaggaagg tggtctctac aatgccatc attgcgataa aggaaggct atcattcaag 5460

55

EP 1 020 527 A1

atgcctctgc cgacagtggc cccaaagatg gaccccccacc cacgaggagc atcgtggaaa 5520  
5 aagaagacgt tccaaccacg tcttcaaagc aagtggattg atgtgacatc tccactgacg 5580  
taagggatga cgcacaatcc cactatcctt cgcaagaccc ttctctata taaggaagtt 5640  
catttcattt ggagaggaca cgctgaaatc accagtctct ctctataaat ctatctctct 5700  
10 ctctataacc atggacccag aacgacgccc ggccgacatc cgcctgcca ccgaggcgga 5760  
catgccggcg gtctgcacca tcgtcaacca ctacatcgag acaagcacgg tcaacttccg 5820  
taccgagccg caggaaccgc aggagtggac ggacgacctc gtccgtctgc gggagcgcta 5880  
15 tccctggctc gtcccgagg tggacggcga ggtccggcg atcgcctacg cgggcccctg 5940  
gaaggcacgc aacgcctacg actggacggc cgagtcgacc gtgtacgtct ccccccgca 6000  
ccagcggacg ggactgggct ccacgtctta caccacctg ctgaagtccc tggaggcaca 6060  
20 gggcttcaag agcgtggctc ctgtcatcgg gctgcccaac gacccgagcg tgcgcatgca 6120  
cgaggcgctc ggatatgccc ccccgggcat gctcggggcg gccggcttca agcacgggaa 6180  
ctggcatgac gtgggtttct ggcagctgga cttcagcctg ccggtaccgc cccgtccggt 6240  
25 cctgcccgtc accgagatct gagatcacgc gttctaggat ccccgatga gctaagctag 6300  
ctatatcatc aatttatgta ttacacataa tatcgactc agtctttcat ctacggcaat 6360  
gtaccagctg atataatcag ttattgaaat atttctgaat ttaaacttgc atcaataaat 6420  
30 ttatgttttt gcttggacta taatacctga cttgttattt tatcaataaa tatttaact 6480  
atatttcttt caagatggga attaacatct acaaatigcc tttcttate gaccatgtac 6540  
35 gtatcgcg 6548

<210> 5

40 <211> 6539

<212> DNA

<213> Escherichia coli LE392

45 <220>

<223> Clone: pTS431

50 <400> 5

aattcaagct tgacgtcagg tggcactttt cggggaatg tgcgcggaac ccctatttgt 60

55

ttatTTTTct aaatacattc aaatatgtat ccgctcatga gacaataacc ctgataaatg 120  
 5 ctccaataat attgaaaaag gaagagtatg agtattcaac atttccgtgt cgcccttatt 180  
 cccttttttg cggcattttg ccttccgtgt ttgtctcacc cagaaacgct ggtgaaagta 240  
 aaagatgctg aagatcagtt ggggtgcacga gtgggttaca tcgaactgga tctcaacagc 300  
 10 ggtaagatcc ttgagagttt tcgccccgaa gaacgttttc caatgatgag cactttttaa 360  
 gttctgctat gtggcgcggt attatcccggt attgacgccg ggcaagagca actcggctcg 420  
 cgcatacact attctcagaa tgacttggtt gagtactcac cagtcacaga aaagcatctt 480  
 15 acggatggca tgacagtaag agaattaigc agtgcgccca taacatgag tgataacact 540  
 gcggccaact tacttctgac aacgatcgga ggaccgaagg agctaaccgc tttttgca 600  
 aacatggggg atcatgtaac tcgcttgat cgttgggaac cggagctgaa tgaagccata 660  
 20 ccaaacgacg agcgtgacac cacgatgcct gtagcaatgg caacaacggt gcgcaacta 720  
 ttaactggcg aactacttac tctagcttcc cggcaacaat taatagactg gatggaggcg 780  
 gataaagttg caggaccact tctgcgctcg gcccttccgg ctggctggtt tattgctgat 840  
 25 aaatctggag ccggtgagcg tgggtctcgc ggtatcattg cagcactggg gccagatggt 900  
 aagccctccc gtatcgtagt tatctacagc acggggagtc aggcaactat ggatgaacga 960  
 aatagacaga tcgtgagat aggtgcctca ctgattaagc attgtaact gtcagaccaa 1020  
 30 gtttactcat atatacttta gattgattta aaacttcatt tttaatttaa aaggatctag 1080  
 gtgaagatcc ttttggctc gagtctcatg accaaaatcc cttaacgiga gtttctgtc 1140  
 35 cactgagcgt cagaccccggt agaaaagatc aaaggatctt cttgagatcc ttttttctg 1200  
 cgcgtaatct gctgcttgca aacaaaaaaa ccaccgctac cagcgggtgt ttgtttgccg 1260  
 gatcaagagc taccaactct ttttccgaag gtaactggct tcagcagagc gcagatacca 1320  
 40 aatactgtcc ttctagtta gccgtagtta ggccaccact tcaagaactc tctagcaccg 1380  
 cctacatacc tcgtctgtct aatcctgta ccagtggctg ctgccagtgg cgataagtcg 1440  
 tgtcttaccg ggttggaactc aagacgatag ttaccggata aggcgcagcg gtcgggctga 1500  
 45 acgggggggt cgtgcacaca gccagcttg gagcgaacga cctacaccga actgagatac 1560  
 ctacagcgtg agcatigaga aagcgccacg ctccccgaag ggagaaaggc ggacaggat 1620  
 ccggtaaagc gcagggtcgg aacaggagag cgcacgaggg agcttcagg gggaaacgcc 1680  
 50 tggatatctt atagtctgt cgggtttcgc cacctctgac ttgagcgtcg atttttgtga 1740  
 tgctcgtcag gggggcggag cctatggaaa aacgccagca acgcggcctt ttacggttc 1800

EP 1 020 527 A1

ctggcctttt gctggccttt tgctcacatg ttctttcctg cgttatcccc tgattctgtg 1860  
5 gataaccgta ttaccgcctt tgagttagct gataccgctc gccgcagccg aacgaccgag 1920  
cgcagcgagt cagttagcga ggaagcgga gagcgccaa tacgcaaacc gcctctcccc 1980  
gcgcgttggc ctgatcagaa ttcttccga tctagtaaca tagatgacac cgcgcgcgat 2040  
10 aatttatcct agtttgcgcg ctatatcttg tttctatcg cgtattaaat gtataattgc 2100  
gggactctaa tcataaaaac ccatctcata aataacgtca tgcatatcat gttaattatt 2160  
acatgcttaa cgtaattcaa cagaaattat atgataatca tcgcaagacc ggcaacagga 2220  
15 ttcaatctta agaaacttta ttgccaaatg ttgaaacgat ctgcttcgga tcctctagat 2280  
tactgatttt tgtaaaggtc tgataatggt ccgttgtttt gtaaatacagc cagtcgcttg 2340  
agtaaagaat ccggtctgaa ttctgaagc ctgagtata gtaatatcc gcttcacgcc 2400  
20 atgttcgtcc gcttttggcc gggagtttgc cttccctgtt tgagaagatg tctccgccga 2460  
tgcttttccc cggagcgacg tctgcaaggt tcccttttga tgccaccag ccgagggctt 2520  
gtgcttctga ttttgtaatg taattatcag gtagcttatg atatgtctga agataatccg 2580  
25 caaccccgctc aaacgtgttg aataaccggt accatcgca cggcttgatg gatctcttgc 2640  
tggacaccgg gatgctagga tgggttatcg tggccggcgt gcgtgtgtgg cttttgtagg 2700  
cgccggcgac ggccggggca atgtggcagg tgagtcacgg tgcaagcgtg cgcaagtac 2760  
30 tgcaacaacc aaggacggtc atggcgaaag cacctcacgc gtccaccgtc tacaggatgt 2820  
agcagtagca cggtgaaaga agtgttgc cgtccattag gtgcattctc accgttggcc 2880  
35 agaacaggac cgttcaacag ttaggttag tgtaggactt ttacgtggtt aatgtatggc 2940  
aaatagtagt aaattttgcc ccattggtc tggctgagat agaacaattt ctggaaagcc 3000  
tctagcatat cttttttgac agctaaactt tgcttcttgc cttcttggtc tagcaatgac 3060  
40 gttgcccatg tctgggcaaa catctggtaa ggtaactgta ttctgttgtt cccttcaacg 3120  
gtcaatccc cacaggccaa gctatcttt ccttggcagt ataggtcctt tgagagatta 3180  
tactaccatt ttttaagtct tataaagacg atgctctcta accagatcga tcagaaacac 3240  
45 aaagtitttag cagcgtaata tccacacac atacacacac gaagctatgc ctctcattt 3300  
tccgagagat tctgacagt accagatgt cagaatgcca ttctatgggc acaagtcgat 3360  
ccacaagctt cttggtggag gtcaagggt gctattatta ttctcttct aggaattat 3420  
50 tcagaattag tgccctttat cataactct ctctgagccg atgtggtttt ggatttcatt 3480  
gttgggagct atgcagttgc ggatattctg ctgtggaaga acaggaaactt atctcgggg 3540

55



EP 1 020 527 A1

gtccttgctg gggcaacatt gatatggttc ctgttcgatg tagtagaata caatataatt 3600  
5 ccgctccttt gccagattgc cattcttgcc atgcttgga tcttcatttg gtcaaagcc 3660  
gcaccactct tggacaggta ttagctttat ttcctgtgga gatggtaga aactcagctt 3720  
acagaaatgg catttcacgt agtataacgc aagacattag gtactaaaac tcaactaact 3780  
10 gtttcgaat ttcagggcc cccaaggat cccagaaatc atcatctctg aacatgcctt 3840  
cagagaaatg gcattgaccg tccattacaa actaacgtac actgtatctg tttttacga 3900  
cattgcatgt ggaaaggatc tgaagagatt tctcctgga cataataatc tactcctttg 3960  
15 ctacgttaat aagagatgta aaaacatgca acagttccag tgccaacatt gtccaaggat 4020  
tgtgcaattc tttctggagc gctaaaattg accagattag acgcatcaga atattgaatt 4080  
gcagagttag ccaataatcc tcataatgtt aatgtgctat tgttggtcac tactcaatat 4140  
20 agttctggac taacaatcag attgtttatg atattaaggt ggttggatct ctattggtat 4200  
tgtcggcgat tggaagttct tgcagcttga caagtctact atatattggt aggtattcca 4260  
gataaatatt aaattttaat aaaacaatca cacagaagga tctgcggcgc ctagcctagg 4320  
25 cccgggccca caaaaatctg agcttaacag cacagttgct cctctcagag cagaatcggg 4380  
tattcaacac cctcatatca actactacgt tgtgtataac ggtccacatg ccggtatata 4440  
cgatgactgg ggttgtacaa aggcggcaac aaacggcgtt cccggagttg cacacaagaa 4500  
30 atttgccact attacagagg caagagcagc agctgacgcg tacacaacaa gtcagcaaac 4560  
agacagggtg aacttcatcc ccaaggaga agctcaactc aagcccaaga gctttgctaa 4620  
ggccctaaca agcccacaa agcaaaaagc cactggctc acgctaggaa ccaaaaggcc 4680  
35 cagcagtgat ccagcccaa aagagatctc ctttgccccg gagattacaa tggacgattt 4740  
cctctatctt tacgatctag gaaggaagt cgaaggtaga ggtgacgaca ctatgttcac 4800  
40 cactgataat gagaaggtaa gcctcttcaa tttcagaaag aatgctgacc cacagatggt 4860  
tagagaggcc tacgcagcag gtctcatcaa gacgatctac ccgagtaaca atctccagga 4920  
gatcaaatac cttccaaga aggttaaga tgcagtcaa agattcagga ctaattgcat 4980  
45 caagaacaca gagaagaca tttttctcaa gatcagaagt actattccag tatggacgat 5040  
tcaaggcttg cttcataaac caaggcaagt aatagagatt ggagtctcta aaaaggtagt 5100  
tcttactgaa tctaaggcca tgcattgagt ctaagattca aatcgaggat ctaacagaac 5160  
50 tcgccgtgaa gactggcgaa cagttcatc agagtctttt acgactcaat gacaagaaga 5220  
aaatcttcgt caacatgggt gagcacgaca ctctggicta ctccaaaaat gtcaaagata 5280

55

5 cagtctcaga agaccaaagg gctattgaga cttttcaaca aaggataatt tcgggaaacc 5340  
 tcctcggatt ccattgccca gctatctgtc acttcatcga aaggacagta gaaaaggaag 5400  
 gtggctccta caaatgccat cattgcgata aaggaaaggc tatcattcaa gatgcctctg 5460  
 ccgacagtgg tcccaaagat ggacccccac ccacgaggag catcgtggaa aaagaagacg 5520  
 10 ttccaaccac gtcttcaaag caagtggatt gatgtgacat ctccactgac gtaagggatg 5580  
 acgcacaatc ccactatcct tcgcaagacc cttctcttat ataaggaagt tcatttcatt 5640  
 tggagaggac acgctgaaat caccagtctc tctctataaa tctatctctc tctctataac 5700  
 15 catggacca gaacgacgcc cggccgacat ccgccgtgcc accgaggcgg acatgccggc 5760  
 ggtctgcacc atcgtcaacc actacatcga gacaagcacg gtcaacttcc gtaccgagcc 5820  
 gcaggaaccg caggagtggc cggacgacct cgtccgtctg cgggagcgct atccctgget 5880  
 20 cgtcgcggag gtggacggcg aggtcgcgg catgcctac gcgggcccct ggaaggcacg 5940  
 caacgcctac gactggacgg ccgagtcgac cgtgtacgtc tcccccgcc accagcggac 6000  
 gggactgggc tccagctct acacccacct gctgaagtcc ctggaggcac agggcttcaa 6060  
 gagcgtggtc gctgtcatcg ggctgcccaa cgacccgagc gtgcgcatgc acgaggcget 6120  
 cggatatgcc ccccgcgga tgcctcgggc ggccggcttc aagcacggga actggcatga 6180  
 30 cgtgggtttc tggcagctgg acttcagcct gccggtaccg ccccgtcgg tctgcccgt 6240  
 caccgagatc tgagatcacg cgttctagga tccccgatg agctaagcta gctatatcat 6300  
 caatttatgt attacacata atatcgcaat cagtcttica tctacggcaa tgtaccagct 6360  
 35 gatataatca gttattgaaa tatttctgaa tttaaacttg catcaataaa tttatgtttt 6420  
 tgcttggact ataatacctg acttggtatt ttatcaataa atatttaaac tatatttctt 6480  
 tcaagatggg aattaacatc tacaattgc cttttcttat cgacatgta cgtatcgcg 6539  
 40

<210> 6

45 <211> 37

<212> DNA

<213> Artificial Sequence

50 <220>

<223> Primer 1

55

<400> 6

5                   cgttcggctc gatggtaccg gttatcaaca cgtttga                   37

10                   <210> 7

                  <211> 38

                  <212> DNA

15                   <213> Artificial Sequence

                  <220>

20                   <223> Primer 2

25                   <400> 7

                  cctctagatt atctgatttt tgtaaaggtc tgataatg                   38

# Claims

- 35 1. A mutant barnase gene having one or some mutations at least in part of the DNA sequence of barnase gene, wherein said mutant barnase gene is capable, when anther-specifically expressed in a plant, of making said plant substantially male sterile without exerting any substantially disadvantageous effect on the tissues except for the anthers.
- 40 2. The gene as claimed in Claim 1, wherein said mutation is a mutation causing frame shift re-coding.
- 45 3. A gene which is derived from a DNA sequence encoding the same amino acid sequence as shown in SEQ ID NO:1 by substitution, deletion, insertion or addition of one to several nucleotides in said DNA sequence, wherein said gene is capable of encoding a protein which makes a plant substantially male sterile when expressed anther-specifically in said plant.
- 50 4. A gene comprising a DNA sequence which is derived from a DNA sequence encoding the same amino acid sequence as shown in SEQ ID NO:1 by substitution, deletion, insertion or addition of one to several nucleotides in the latter DNA sequence, and a promoter located upstream of said DNA sequence for allowing an anther-specific expression, wherein said gene is capable of making a plant substantially male sterile when introduced into the genome of said plant.
- 55 5. A gene represented by SEQ ID NO:3 which encodes a protein capable of making a plant substantially male sterile when expressed anther-specifically in said plant.
6. A gene comprising the sequence represented by SEQ ID NO:3 and a promoter located upstream of said sequence for allowing an anther-specific expression, wherein said gene is capable of making a plant substantially male sterile when introduced into the genome of said plant.

**EP 1 020 527 A1**

7. A recombinant vector which contains a gene as claimed in any of Claims 1 to 6 and expresses said gene in a host plant.
8. A method of making a plant male sterile which comprises transforming said plant by a mutant barnase gene as claimed in any of Claims 1 to 6 and allowing said mutant barnase gene to be expressed anther-specifically.
9. The method as claimed in Claim 8, wherein said plant is transformed by said mutant barnase gene by integrating said gene into the genome of said plant.
10. A transgenic plant wherein a gene as claimed in any of Claims 1 to 6 has been introduced.

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP99/04167

## A. CLASSIFICATION OF SUBJECT MATTER

Int.Cl<sup>6</sup> C12N15/55, C12N9/22, C12N15/82, A01H1/00, A01H5/00

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

Int.Cl<sup>6</sup> C12N15/55, C12N9/22, C12N15/82, A01H1/00, A01H5/00

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)  
WPI/BIOSIS (DIALOG), REGISTRY (STN)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO, 96/26283, A (Plant Genetic Systems N.V.), 29 August, 1996 (29. 08. 96) & EP, 811070, A1 & JP, 11-500617, A	1-10
A	Journal of Immunology Vol. 147 No. 8 (1991) Fettes J.V. et al., "A frameshift mutation at the amino terminus of the nucleoprotein gene does not affect generation of cytotoxic T lymphocyte epitopes" p.2697-2705	1-10

☐ Further documents are listed in the continuation of Box C.☐ See patent family annex.

\* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, etc, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"d" document member of the same patent family

Date of the actual completion of the international search  
28 October, 1999 (28. 10. 99)Date of mailing of the international search report  
9 November, 1999 (09. 11. 99)Name and mailing address of the ISA/  
Japanese Patent Office

Authorized officer

Facsimile No.

Telephone No.

Form PCT/ISA/210 (second sheet) (July 1992)

